

APCalcBC-HomeworkQuiz-#4

1. If $y = \arctan(\cos x)$, then $\frac{dy}{dx} =$
- (A) $\frac{-\sin x}{1+\cos^2 x}$
 (B) $-(\text{arcsec}(\cos x))^2 \sin x$
 (C) $(\text{arcsec}(\cos x))^2$
 (D) $\frac{1}{(\arccos x)^2 + 1}$
 (E) $\frac{1}{1+\cos^2 x}$
- $\frac{1}{1+(\cos x)^2} (-\sin x)$
2. $\int x^2 \cos(x^3) dx =$
- (A) $-\frac{1}{3} \sin(x^3) + C$
 (B) $\frac{1}{3} \sin(x^3) + C$
 (C) $-\frac{x^3}{3} \sin(x^3) + C$
 (D) $\frac{x^3}{3} \sin(x^3) + C$
 (E) $\frac{x^3}{3} \sin\left(\frac{x^4}{4}\right) + C$
- $\underline{u\text{-sub}}$ $u = x^3$
 $\frac{du}{dx} = 3x^2$
 $du = 3x^2 dx$
 $x^2 dx = \frac{1}{3} du$
- $\frac{1}{3} \int \cos(u) du$
 $\frac{1}{3} \sin u + C$
 $\frac{1}{3} \sin(x^3) + C$
3. $\lim_{x \rightarrow \infty} \frac{x^3 - 2x^2 + 3x - 4}{4x^3 - 3x^2 + 2x - 1} =$
- (A) 4
 (B) 1
 (C) 1/4
 (D) 0
 (E) -1
- ∞ multiple L'Hopital or $\frac{x^3 - 2x^2 + 3x - 4}{4x^3 - 3x^2 + 2x - 1} = \frac{1}{4}$
4. The third-degree Taylor polynomial about $x = 0$ of $\ln(1-x)$ is
- (A) $-x - \frac{x^2}{2} - \frac{x^3}{3}$
 (B) $1 - x + \frac{x^2}{2}$
 (C) $x - \frac{x^2}{2} + \frac{x^3}{3}$
 (D) $-1 + x - \frac{x^2}{2}$
 (E) $-x + \frac{x^2}{2} - \frac{x^3}{3}$
- $P_3 = f(0) + f'(0)(x-0) + \frac{f''(0)}{2!}(x-0)^2 + \frac{f'''(0)}{3!}(x-0)^3$
 $f(x) = \ln(1-x) \quad f(0) = \ln(1) = 0$
 $f'(x) = \frac{1}{1-x}(-1) \quad f'(0) = \frac{1}{1-0} = -1$
 $f''(x) = \frac{1}{(1-x)^2}(-1) = \frac{-1}{(1-x)^2} \quad f''(0) = -1$
 $f'''(x) = \frac{0 - (-1)(2(1-x)(-1))}{(1-x)^4} = \frac{-2}{(1-x)^3} \quad f'''(0) = -2$
- $P_3 = 0 + (-1)x + \frac{(-1)}{2}x^2 + \frac{(-2)}{3!}x^3 = -x - \frac{1}{2}x^2 - \frac{1}{3}x^3$
5. If $y = \frac{\ln x}{x}$, then $\frac{dy}{dx} =$

APCalcBC-HomeworkQuiz #4

$$y = \frac{\ln x}{x} \quad \frac{dy}{dx} = \frac{x(\frac{1}{x}) - \ln x(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- (A) $\frac{1}{x}$
 (B) $\frac{1}{x^2}$
 (C) $\frac{\ln x - 1}{x^2}$
 (D) $\frac{1 - \ln x}{x^2}$
 (E) $\frac{1 + \ln x}{x^2}$

6. A pizza, heated to a temperature of 350 degrees Fahrenheit (${}^{\circ}\text{F}$) is taken out of an oven and placed in a (75°F) room at time $t = 0$ minutes. The temperature of the pizza is changing at a rate of $-110e^{-0.4t}$ degrees Fahrenheit per minute. To the nearest degree, what is the temperature of the pizza at time $t = 5$ minutes?

- (A) 112°F
 (B) 119°F
 (C) 147°F
 (D) 238°F
 (E) 335°F

$$\int_0^5 (-110e^{-0.4t}) dt = T(5) - T(0)$$

$$-237.282721 = T(5) - 350$$

$$T(5) = 112.217^{\circ}\text{F}$$

7. $\int \frac{7x}{(2x-3)(x+2)} dx =$

Partial fraction expansion: $\frac{7x}{(2x-3)(x+2)} = \frac{A}{2x-3} + \frac{B}{x+2}$

$$A(x+2) + B(2x-3) = 7x$$

$$(A+2B)x + (2A-3B) = (7)x + b$$

- (A) $\frac{3}{2} \ln|2x-3| + 2 \ln|x+2| + C$
 (B) $3 \ln|2x-3| + 2 \ln|x+2| + C$
 (C) $3 \ln|2x-3| - 2 \ln|x+2| + C$
 (D) $-\frac{6}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$
 (E) $-\frac{3}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$

$$3 \left\{ \int \frac{1}{2x-3} dx + 2 \int \frac{1}{x+2} dx \right.$$

$$\begin{aligned} u &= 2x-3 \\ du &= 2dx, dx = \frac{1}{2}du \end{aligned} \quad \begin{aligned} u &= x+2 \\ du &= dx \end{aligned}$$

$$\frac{3}{2} \int \frac{1}{u} du + 2 \int \frac{1}{u} du$$

$$\begin{cases} A+2B=7 \\ 2A-3B=0 \end{cases}$$

$$\begin{cases} -4A-4B=-14 \\ 2A-3B=0 \end{cases}$$

$$\begin{cases} -7B=-14 \\ B=2 \end{cases}$$

$$\begin{cases} A+2(2)=7 \\ A=3 \end{cases}$$

| | | | | | |
|------|---|----|----|---|----|
| x | 2 | 3 | 5 | 8 | 13 |
| f(x) | 6 | -2 | -1 | 3 | 9 |

The function f is continuous on the closed interval $[2, 13]$ and has values as shown in the table above. Using the

intervals $[2, 3]$, $[3, 5]$, $[5, 8]$, and $[8, 13]$ what is the approximation of $\int_2^{13} f(x) dx$ obtained from a left

Riemann sum?

| | interval | x_L | $f(x_L) \cdot \Delta x = \text{area}$ |
|--------|-----------|-------|---------------------------------------|
| (A) 6 | $[2, 3]$ | 2 | $6 \cdot 1 = 6$ |
| (B) 14 | $[3, 5]$ | 3 | $-2 \cdot 2 = -4$ |
| (C) 28 | $[5, 8]$ | 5 | $-1 \cdot 3 = -3$ |
| (D) 32 | $[8, 13]$ | 8 | $3 \cdot 5 = 15$ |
| (E) 50 | | | |

$$\frac{-12}{14}$$


APCalcBC-HomeworkQuiz-#4

9.

(CALCULATOR ALLOWED)

What is the area enclosed by the curves $y = x^3 - 8x^2 + 18x - 5$ and $y = x + 5$?

- (A) 10.667
 (B) 11.833
 (C) 14.583
 (D) 21.333
 (E) 32




$$A = \int_{1}^{5} [(x^3 - 8x^2 + 18x - 5) - (x + 5)] dx \quad 58.33$$

$$+ \int_{1}^{5} [(x + 5) - (x^3 - 8x^2 + 18x - 5)] dx \quad 11.25$$

$$= 11.833$$

10. The top of a 25-foot ladder is sliding down a vertical wall at a constant rate of 3 feet per minute. When the top of the ladder is 7 feet from the ground, what is the rate of change of the distance between the bottom of the ladder and the wall?

- (A) $-\frac{7}{8}$ feet per minute
 (B) $-\frac{7}{24}$ feet per minute
 (C) $\frac{7}{24}$ feet per minute
 (D) $\frac{7}{8}$ feet per minute
 (E) $\frac{21}{25}$ feet per minute



$$x^2 + y^2 = 25^2$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$2(25) \frac{dx}{dt} + 2(7)(-3) = 0$$

$$\frac{dx}{dt} = \frac{42}{48} = \frac{7}{8} \text{ ft/min}$$

11. The length of the path described by the parametric equations $x = \cos^3 t$ and $y = \sin^3 t$, for $0 \leq t \leq \frac{\pi}{2}$ is given by

- (A) $\int_0^{\frac{\pi}{2}} \sqrt{3\cos^2 t + 3\sin^2 t} dt$
 (B) $\int_0^{\frac{\pi}{2}} \sqrt{-3\cos^2 t \sin t + 3\sin^2 t \cos t} dt$
 (C) $\int_0^{\frac{\pi}{2}} \sqrt{9\cos^4 t + 9\sin^4 t} dt$
 (D) $\int_0^{\frac{\pi}{2}} \sqrt{9\cos^4 t \sin^2 t + 9\sin^4 t \cos^2 t} dt$
 (E) $\int_0^{\frac{\pi}{2}} \sqrt{\cos^6 t + \sin^6 t} dt$

$$\text{arc length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$= \int_0^{\frac{\pi}{2}} \sqrt{(3\cos^2 t(-\sin t))^2 + (3\sin^2 t \cos t)^2} dt$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} \sqrt{9\cos^4 t \sin^2 t + 9\sin^4 t \cos^2 t} dt$$

$$= \int_0^{\frac{\pi}{2}} \sqrt{9(\sin^2 t + \cos^2 t)(\cos^4 t + \sin^4 t)} dt$$

$$= \int_0^{\frac{\pi}{2}} 3 \sqrt{\cos^4 t + \sin^4 t} dt$$

APCalcBC-HomeworkQuiz-#4

12. Which of the following series diverge?



I. $\sum_{n=0}^{\infty} \left(\frac{\sin 2}{\pi}\right)^n$ geometric $r = \frac{\sin 2}{\pi} \approx \frac{0.9}{3.14} < 1$, converges


II. $\sum_{n=1}^{\infty} \frac{1}{\sqrt[3]{n}}$ p-series $\sum_{n=1}^{\infty} \frac{1}{n^{1/3}}$ w/p = 1/3, diverges

III. $\sum_{n=1}^{\infty} \left(\frac{e^n}{e^n + 1}\right)$ nth term test $\lim_{n \rightarrow \infty} \frac{e^n}{e^n + 1} = 1 \neq 0$, diverges

- (A) III only
 (B) I and II only
 (C) I and III only
 (D) II and III only
 (E) I, II, and III

13. The region R in the first quadrant is enclosed by the lines $x=0$ and $y=5$ and the graph of $y = x^2 + 1$. The volume of the solid generated when R is revolved about the y-axis is

- (A) 6π
 (B) 8π
 (C) $34\pi/3$
 (D) 16π
 (E) $544\pi/15$



disc $V = \int \pi r^2 dy$ $x^2 = y - 1$
 $= \int \pi (\sqrt{y-1})^2 dy$
 $= \pi \int_1^5 (y-1) dy = \pi \left(\frac{1}{2}y^2 - y \right) \Big|_1^5$
 $= \pi \left(\frac{25}{2} - 5 \right) - \pi \left(\frac{1}{2} - 1 \right) = \pi \left(\frac{25}{2} - \frac{1}{2} \right) = \pi(12) = 12\pi$

14. A curve is described by the parametric equations $x = t^2 + 2t$ and $y = t^3 + t^2$. An equation of the line tangent to the curve at the point determined by $t = 1$ is

- (A) $2x - 3y = 0$
 (B) $4x - 5y = 2$
 (C) $4x - y = 10$
 (D) $5x - 4y = 7$
 (E) $5x - y = 13$

$(x, y) = (3, 2)$
 $m = \frac{dy}{dx} = \frac{(dy/dt)}{(dx/dt)} = \frac{3t^2 + 2t}{2t + 2} \Big|_{(t=1)} = \frac{5}{4}$


$(y - 2) = \frac{5}{4}(x - 3)$

$4y - 8 = 5x - 15$

$5x - 4y = 7$

APCalcBC-HomeworkQuiz-#4

15.



The graph of a function f is shown above. At which value of x is f continuous, but not differentiable?

- (A) a
 (B) b
 (C) c
 (D) d
 (E) e

16. $\int_1^\infty \frac{x}{(1+x^2)^2} dx$ is

$$= \lim_{b \rightarrow \infty} \int_1^b \frac{x}{(1+x^2)^2} dx$$

- (A) $-\frac{1}{2}$
 (B) $-\frac{1}{4}$
 (C) $\frac{1}{4}$
 (D) $\frac{1}{2}$
 (E) divergent

$$u\text{-sub: } u = 1+x^2$$

$$\frac{du}{dx} = 2x$$

$$dx = \frac{1}{2x} du, x dx = \frac{1}{2} du$$

$$\frac{1}{2} \int u^{-2} du, \frac{1}{2} \cdot \frac{u^{-1}}{-1} = \frac{-1}{2(u+1^2)}$$

$$\lim_{b \rightarrow \infty} \left[\frac{-1}{2(1+b^2)} \right] - \left[\frac{-1}{2(1+1^2)} \right]$$

$$0 + \frac{1}{4} = \frac{1}{4}$$

17. If $y = \cos^2 3x$, then $dy/dx =$

- (A) $-6 \sin 3x \cos 3x$
 (B) $-2 \cos 3x$
 (C) $2 \cos 3x$
 (D) $6 \cos 3x$
 (E) $2 \sin 3x \cos 3x$

$$y = (\cos(3x))^2$$


$$\frac{dy}{dx} = 2(\cos(3x))'(-\sin(3x)) \cdot 3$$

$$= -6 \sin(3x) \cos(3x)$$

18. A rectangular area is to be enclosed by a wall on one side and fencing on the other three sides. If 18 meters of fencing are used, what is the maximum area that can be enclosed?

APCalcBC-HomeworkQuiz-#4

- (A) $\frac{9}{2} m^2$
 (B) $\frac{81}{4} m^2$
 (C) $27 m^2$
 (D) $40 m^2$
 (E) $\frac{81}{2} m^2$

constraint

$$F = x + 2y = 18$$

$$x = 18 - 2y$$


$$A = 18y - 2y^2$$

$$A' = 18 - 4y = 0$$

$$4y = 18, y = \frac{18}{4} = \frac{9}{2}, x = 18 - 2\left(\frac{9}{2}\right) = 9$$

$$Area = \frac{81}{2} m^2$$

19.

Graph of f'

The graph of f' , the derivative of the function f is shown above. On which of the following intervals is f decreasing? when $f' < 0$ ($0, 2 \cup 4, 6$)

- (A) $[2, 4]$ only
 (B) $[3, 5]$ only
 (C) $[0, 1]$ and $[3, 5]$
 (D) $[2, 4]$ and $[6, 7]$
 (E) $[0, 2]$ and $[4, 6]$ (nearly should be open intervals)

20. The velocity vector of a particle moving in the xy -plane has components given by $\frac{dx}{dt} = \sin(t^2)$ and $\frac{dy}{dt} = e^{\cos t}$. At time $t = 4$, the position of the particle is $(2, 1)$. What is the y-coordinate of the position vector at time $t = 3$?

- (A) 0.410
 (B) 0.590
 (C) 0.851
 (D) 1.410

$$\int_3^4 e^{\cos t} dt = y(4) - y(3)$$

$$0.4097023586 = 1 - y(3)$$

$$y(3) = 1 - 0.4097023586 = 0.59029$$

21. $\int xe^{2x} dx =$

- (A) $xe^{2x}/2 - e^{2x}/4 + C$
 (B) $xe^{2x}/2 - e^{2x}/2 + C$
 (C) $xe^{2x}/2 + e^{2x}/4 + C$
 (D) $xe^{2x}/2 + e^{2x}/2 + C$
 (E) $x^2e^{2x}/4 + C$

By parts: $u = x, dv = e^{2x} dx, uv - \int v du$

$$\frac{du}{dt} = 1, \int dv = \int e^{2x} dx, \frac{1}{2}xe^{2x} - \frac{1}{2}\int e^{2x} dx$$

$$du = dx, v = \frac{1}{2}e^{2x}, \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + C$$

APCalcBC-HomeworkQuiz #4

22. What is the general solution to the differential equation $\frac{dy}{dx} = \frac{x \cos(x^2)}{4y}$ for $y > 0$?

- (A) $y = \frac{1}{2} \sqrt{\sin(x^2)} + C$
 (B) $y = \sqrt{\frac{1}{4} \sin(x^2)} + C$
 (C) $y = \frac{1}{8} \sin(x^2) + C$
 (D) $y = Ce^{\frac{1}{8} \sin(x^2)}$

$$\int 4y \, dy = \int x \cos(x^2) \, dx$$

$$\begin{aligned} u &= x^2 \\ du &= 2x \end{aligned}$$

$$\begin{aligned} 2y^2 &= \frac{1}{2} \int \cos(u) \, du \\ 2y^2 &= \frac{1}{2} \sin(u) + C \\ y^2 &= \frac{1}{4} \sin(x^2) + C \\ y &= \pm \sqrt{\frac{1}{4} \sin(x^2) + C} \end{aligned}$$

23. Let f be a differentiable function such that $f'(x) \geq 1$ for all x . If $a < b$, which of the following statements could be false?

- (A) $\frac{f(b)-f(a)}{b-a} \geq 1$ true
 (B) $f(b) > f(a)$ true
 (C) There is a value c in the open interval (a, b) such that $f(c) = 0$. ($f(a)$ & $f(b)$ might both be positive or both negative)
 (D) There is a value c in the open interval (a, b) such that $f(c) = \frac{f(a)+f(b)}{2}$. guaranteed by I.V.T.

24. The Taylor series for $\ln x$, centered at $x=1$, is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$. Let f be the function given by the sum of the

first three nonzero terms of this series. The maximum value of $|\ln x - f(x)|$ for $0.3 \leq x \leq 1.7$ is

- (A) 0.030 $f(x) = \ln(x) = \frac{1}{n} x^n$
 (B) 0.039 $f'(x) = \frac{1}{x} = x^{-1}$
 (C) 0.145 $f''(x) = -\frac{1}{x^2} = -\frac{1}{x^2}$
 (D) 0.153 $f'''(x) = 2x^{-3} = \frac{2}{x^3}$
 (E) 0.529 error = $\max_{n=3} f''(x) \text{ in interval } (x-1)^n = \frac{(0.3)^3}{4!} (0.3-1)^4$
 for $n=3$

25. For what value of k , if any, will $y = ke^{-2x} + 4 \cos(3x)$ be a solution to the differential equation $y'' + 9y = 26e^{-2x}$?

- (A) 2
 (B) $\frac{13}{5}$
 (C) 26
 (D) There is no such value of k .

$$y' = -2ke^{-2x} - 12 \sin(3x)$$

$$y'' = 4ke^{-2x} - 36 \cos(3x)$$

$$[4ke^{-2x} - 36 \cos(3x)] + 9[ke^{-2x} + 4 \cos(3x)] = 26e^{-2x}$$

$$(4k+9k)e^{-2x} = 26e^{-2x}$$

$$13k = 26$$

$$k = 2$$


this is also an alternating series so error should be less than 1st neglected term; $-\frac{1}{4}(x-1)^4$ which over interval is no larger than $\frac{1}{4}(1.7)^4 = .060025$, also not helpful

* However, if in a calculator you enter:

$$y_1 = (x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3$$

$$y_2 = \ln x$$

$$y_3 = y_2 - y_1 \text{ (the error) and just graph } y_3 :$$



trace, value at x=.03, error = -.1444