


2005 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS



2. The curve above is drawn in the xy -plane and is described by the equation in polar coordinates $r = \theta + \sin(2\theta)$ for $0 \leq \theta \leq \pi$, where r is measured in meters and θ is measured in radians. The derivative of r with respect to θ is given by $\frac{dr}{d\theta} = 1 + 2\cos(2\theta)$.
- (a) Find the area bounded by the curve and the x -axis.
 - (b) Find the angle θ that corresponds to the point on the curve with x -coordinate -2 .
 - (c) For $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$, $\frac{dr}{d\theta}$ is negative. What does this fact say about r ? What does this fact say about the curve?
 - (d) Find the value of θ in the interval $0 \leq \theta \leq \frac{\pi}{2}$ that corresponds to the point on the curve in the first quadrant with greatest distance from the origin. Justify your answer.

WRITE ALL WORK IN THE TEST BOOKLET.

2005 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

Distance x (cm)	0	1	5	6	8
Temperature $T(x)$ (°C)	100	93	70	62	55

3. A metal wire of length 8 centimeters (cm) is heated at one end. The table above gives selected values of the temperature $T(x)$, in degrees Celsius (°C), of the wire x cm from the heated end. The function T is decreasing and twice differentiable.
- (a) Estimate $T'(7)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Write an integral expression in terms of $T(x)$ for the average temperature of the wire. Estimate the average temperature of the wire using a trapezoidal sum with the four subintervals indicated by the data in the table. Indicate units of measure.
- (c) Find $\int_0^8 T'(x) dx$, and indicate units of measure. Explain the meaning of $\int_0^8 T'(x) dx$ in terms of the temperature of the wire.
- (d) Are the data in the table consistent with the assertion that $T''(x) > 0$ for every x in the interval $0 < x < 8$? Explain your answer.
-

WRITE ALL WORK IN THE TEST BOOKLET.

END OF PART A OF SECTION II

2005 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

**CALCULUS BC
SECTION II, Part B**

Time—45 minutes


Number of problems—3

No calculator is allowed for these problems.

4. Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated, and sketch the solution curve that passes through the point $(0, 1)$.


(Note: Use the axes provided in the pink test booklet.)



- (b) The solution curve that passes through the point $(0, 1)$ has a local minimum at $x = \ln\left(\frac{3}{2}\right)$. What is the y -coordinate of this local minimum?
- (c) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(0) = 1$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $f(-0.4)$. Show the work that leads to your answer.
- (d) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine whether the approximation found in part (c) is less than or greater than $f(-0.4)$. Explain your reasoning.

WRITE ALL WORK IN THE TEST BOOKLET.

2005 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS



5. A car is traveling on a straight road. For $0 \leq t \leq 24$ seconds, the car's velocity $v(t)$, in meters per second, is modeled by the piecewise-linear function defined by the graph above.
- (a) Find $\int_0^{24} v(t) dt$. Using correct units, explain the meaning of $\int_0^{24} v(t) dt$.
- (b) For each of $v'(4)$ and $v'(20)$, find the value or explain why it does not exist. Indicate units of measure.
- (c) Let $a(t)$ be the car's acceleration at time t , in meters per second per second. For $0 < t < 24$, write a piecewise-defined function for $a(t)$.
- (d) Find the average rate of change of v over the interval $8 \leq t \leq 20$. Does the Mean Value Theorem guarantee a value of c , for $8 < c < 20$, such that $v'(c)$ is equal to this average rate of change? Why or why not?
-

6. Let f be a function with derivatives of all orders and for which $f(2) = 7$. When n is odd, the n th derivative of f at $x = 2$ is 0. When n is even and $n \geq 2$, the n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n-1)!}{3^n}$.
- (a) Write the sixth-degree Taylor polynomial for f about $x = 2$.
- (b) In the Taylor series for f about $x = 2$, what is the coefficient of $(x-2)^{2n}$ for $n \geq 1$?
- (c) Find the interval of convergence of the Taylor series for f about $x = 2$. Show the work that leads to your answer.
-

WRITE ALL WORK IN THE TEST BOOKLET.

END OF EXAM

2006 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

3. An object moving along a curve in the xy -plane is at position $(x(t), y(t))$ at time t , where

$$\frac{dx}{dt} = \sin^{-1}(1 - 2e^{-t}) \text{ and } \frac{dy}{dt} = \frac{4t}{1 + t^3}$$

for $t \geq 0$. At time $t = 2$, the object is at the point $(6, -3)$. (Note: $\sin^{-1} x = \arcsin x$)

- (a) Find the acceleration vector and the speed of the object at time $t = 2$.
 - (b) The curve has a vertical tangent line at one point. At what time t is the object at this point?
 - (c) Let $m(t)$ denote the slope of the line tangent to the curve at the point $(x(t), y(t))$. Write an expression for $m(t)$ in terms of t and use it to evaluate $\lim_{t \rightarrow \infty} m(t)$.
 - (d) The graph of the curve has a horizontal asymptote $y = c$. Write, but do not evaluate, an expression involving an improper integral that represents this value c .
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

END OF PART A OF SECTION II


2002 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

CALCULUS BC
SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.



Graph of f

4. The graph of the function f shown above consists of two line segments. Let g be the function given by $g(x) = \int_0^x f(t) dt$.
- (a) Find $g(-1)$, $g'(-1)$, and $g''(-1)$.
 - (b) For what values of x in the open interval $(-2, 2)$ is g increasing? Explain your reasoning.
 - (c) For what values of x in the open interval $(-2, 2)$ is the graph of g concave down? Explain your reasoning.
 - (d) On the axes provided, sketch the graph of g on the closed interval $[-2, 2]$.
(Note: The axes are provided in the pink test booklet only.)