

Sample Examination II

Section I Part A

Directions: Solve each of the following problems, using available space for scratchwork. After examining the form of the choices, decide which is the best of the choices given. Do not spend too much time on any one problem. Calculators may NOT be used on this part of the exam.

In this test: Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.

skip #2 (2nd semester topic)

Part I - M/C calculator not allowed

1. $\int_0^2 (2x^3 + 3) dx =$

(A) 8

(B) 11

(C) 14

(D) 20

(E) 24

CalcAB Sample Exam II (online version)

2. The slope field for the differential equation $\frac{dy}{dx} = \frac{3y}{xy + 5x}$ will have vertical segments when

- (A) $x = 0$, only
- (B) $y = 0$, only
- (C) $y = -5$, only
- (D) $y = 5$, only
- (E) $x = 0$ or $y = -5$

SKIP

3. Suppose that f is a continuous function defined for all real numbers x and $f(-5) = 3$ and $f(-1) = -2$. If $f(x) = 0$ for one and only one value of x , then which of the following could be x ?

- (A) -7
- (B) -2
- (C) 0
- (D) 1
- (E) 2

CalcAB Sample Exam II (online version)

4. If $f(x) = (2 + 3x)^4$, then the fourth derivative of f is

- (A) 0
- (B) $4!(3)$
- (C) $4!(3^4)$
- (D) $4!(3^5)$
- (E) $4!(2 + 3x)$

5. At what value(s) of x does $f(x) = x^4 - 8x^2$ have a relative minimum?

- (A) 0 and -2 only
- (B) 0 and 2 only
- (C) 0 only
- (D) -2 and 2 only
- (E) -2 , 0, and 2

6. $\int \sqrt{x}(x+2) dx =$

(A) $\sqrt{x^3} + 2\sqrt{x} + C$ (B) $\frac{2}{5}x^{\frac{3}{2}} + \frac{4}{3}x^{\frac{1}{2}} + C$ (C) $\frac{3}{2}\sqrt{x} + \frac{1}{\sqrt{x}} + C$

(D) $\frac{2}{5}x^{\frac{5}{2}} + \frac{4}{3}x^{\frac{3}{2}} + C$ (E) $\frac{x^2}{2} \left(\frac{2}{3}x^{\frac{3}{2}} + 2x \right) + C$

7. The $\lim_{h \rightarrow 0} \frac{|x+h| - |x|}{h}$ at $x = 3$ is

(A) -1

(B) 0

(C) 1

(D) 3

(E) nonexistent

CalcAB Sample Exam II (online version)

8. The function $y = x^4 + bx^2 + 8x + 1$ has a horizontal tangent and a point of inflection for the same value of x . What must be the value of b ?

(A) -6

(B) -1

(C) 1

(D) 4

(E) 6

9. If $y = 7$ is a horizontal asymptote of a rational function f , then which of the following must be true?

(A) $\lim_{x \rightarrow 7} f(x) = \infty$

(B) $\lim_{x \rightarrow -\infty} f(x) = -7$

(C) $\lim_{x \rightarrow 0} f(x) = 7$

(D) $\lim_{x \rightarrow 7} f(x) = 0$

(E) $\lim_{x \rightarrow \infty} f(x) = 7$

CalcAB Sample Exam II (online version)

10. Let $f(x) = e^{(x^2)}$. At how many points in the interval $[-a, a]$, does the instantaneous rate of change of f equal the average rate of change of f ?


- (A) None
- (B) One
- (C) Two
- (D) Three
- (E) More than three

11. Let f be the function given by $f(x) = x^3$. What are all values of c that satisfy the conclusion of the Mean Value Theorem on the closed interval $[-1, 2]$?

- (A) 0 only
- (B) 1 only
- (C) $\sqrt{3}$ only
- (D) -1 and 1
- (E) $-\sqrt{3}$ and $\sqrt{3}$

CalcAB Sample Exam II (online version)

12.



For the figure above, the area of the shaded region is $\ln 4$ when k is

- (A) 4
- (B) 8
- (C) e
- (D) e^2
- (E) e^3

13. If $x + y = xy$, then $\frac{dy}{dx}$ is

(A) $\frac{1}{x-1}$

(B) $\frac{y-1}{x-1}$

(C) $\frac{1-y}{x-1}$

(D) $x + y - 1$

(E) $\frac{2-xy}{y}$

CalcAB Sample Exam II (online version)

4. If f and g are continuously differentiable functions defined for all real numbers, which of the following definite integrals is equal to $f(g(4)) - f(g(2))$?

(A) $\int_2^4 f'(g(x)) \, dx$

(B) $\int_2^4 f(g(x))f'(x) \, dx$

(C) $\int_2^4 f(g(x))g'(x) \, dx$

(D) $\int_2^4 f'(g(x))g'(x) \, dx$

(E) $\int_2^4 f(g'(x))g'(x) \, dx$

15. The velocity of a particle moving along the y -axis is given by $v(t) = 8 - 2t$ for $t \geq 0$. The particle moves upward until it reaches the origin and then moves downward. The position of the particle at any time t is given by

(A) $-t^2 + 8t - 16$

(B) $-t^2 + 8t + 16$

(C) $2t^2 - 8t - 16$

(D) $8t - 2t^2$

(E) $8t - t^2$

CalcAB Sample Exam II (online version)

16. If the substitution $u = \sqrt{x-1}$ is made, the integral $\int_2^5 \frac{\sqrt{x-1}}{x} dx =$

(A) $\int_2^5 \frac{2u^2}{u^2+1} du$

(B) $\int_1^2 \frac{u^2}{u^2+1} du$

(C) $\int_1^2 \frac{u^2}{2(u^2+1)} du$

(D) $\int_2^5 \frac{u}{u^2+1} du$

(E) $\int_1^2 \frac{2u^2}{u^2+1} du$

17. What are all values of x for which the function $f(x) = x^3 + 6x^2 + 9x + 1$ is increasing?

(A) $(-\infty, -3)$ only

(B) $(-3, -1)$ only

(C) $(-1, \infty)$ only

(D) $(-\infty, -3) \cup (-1, \infty)$

(E) $(-\infty, -3) \cup (1, \infty)$

CalcAB Sample Exam II (online version)

18. If $\int_0^2 (2x^3 - kx^2 + 2k) dx = 12$, then k must be

(A) -3

(B) -2

(C) 1

(D) 2

(E) 3

19. $\int (\sec^2 x)(\tan^2 x) dx =$

(A) $\frac{1}{3} \tan^3 x + C$

(B) $\tan^3 x + C$

(C) $\frac{1}{2} \tan^2 x + C$

(D) $\frac{1}{3} \sec^3 x + C$

(E) $\tan^2 x + C$

CalcAB Sample Exam II (online version)

20. For $|x| < 1$, the derivative of $y = \ln \sqrt{1 - x^2}$ is

- (A) $\frac{x}{1 - x^2}$ (B) $\frac{x}{x^2 - 1}$ (C) $\frac{-1}{x^2 - 1}$ (D) $\frac{1}{2(1 - x^2)}$ (E) $\frac{1}{\sqrt{1 - x^2}}$

21. A function whose derivative is a constant multiple of itself must be

- (A) periodic
(B) linear
(C) exponential
(D) quadratic
(E) logarithmic

CalcAB Sample Exam II (online version)

23. The edge of a cube is increasing at the rate of 0.05 centimeters per second. In terms of the edge of the cube, s , what is the rate of change of the volume of the cube, in cubic centimeters per second?

(A) 0.05^3

(B) $0.05s^2$

(C) $0.05s^3$

(D) $0.15s^2$

(E) $3s^2$

24. The tangent line to the graph $y = e^{2-x}$ at the point $(1, e)$ intersects both coordinate axes. What is the area of the triangle formed by this tangent line and the coordinate axes?

(A) $2e$

(B) $e^2 - 1$

(C) e^2

(D) $2e\sqrt{e}$

(E) $4e$

CalcAB Sample Exam II (online version)

25. If f is the function defined by $f(x) = \frac{5x^7}{7} + 4x^6 + 6x^5 + x + 1$, what are all the x -coordinates of the points of inflection of the graph of f ?






- (A) -2 only
- (B) 0 only
- (C) 2 only
- (D) -2 and 0 only
- (E) $-2, 0,$ and 2

26. A normal line to the graph of a function f at the point $(x, f(x))$ is defined to be the line perpendicular to the tangent line at that point. The equation of the normal line to the curve $y = \sqrt[3]{x^2 - 1}$ at the point where $x = 3$ is

- (A) $y + 12x = 38$
- (B) $y - 4x = 10$
- (C) $y + 2x = 4$
- (D) $y + 2x = 8$
- (E) $y - 2x = -4$

CalcAB Sample Exam II (online version)

27. Which graph best represents the position of a particle, $s(t)$, as a function of time, if the particle's velocity and acceleration are both positive?



Answer

28. If n is a positive integer, then $\lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(\frac{1}{n}\right)^2 + \left(\frac{2}{n}\right)^2 + \dots + \left(\frac{n-1}{n}\right)^2 \right] =$

(A) $\int_0^1 \frac{1}{x^2} dx$

(B) $\int_0^1 x^2 dx$

(C) $\int_0^1 \frac{2}{x^2} dx$

(D) $\int_0^1 \frac{1}{x} dx$

(E) $\int_0^2 x^2 dx$


CalcAB Sample Exam II (online version)

Part II - M/C calculator allowed

29. If f is a function such that $\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = 0$, which of the following must be true?

- (A) $\lim_{x \rightarrow a} f(x)$ does not exist
- (B) $f(a)$ does not exist
- (C) $f'(a) = 0$
- (D) $f(a) = 0$
- (E) $f(x)$ is continuous at $x = 0$

30.



The graph of the derivative of a twice-differentiable function f is shown above. If $f(1) = -2$, which of the following is true?

- (A) $f(2) < f'(2) < f''(2)$
- (B) $f''(2) < f'(2) < f(2)$
- (C) $f'(2) < f(2) < f''(2)$
- (D) $f(2) < f''(2) < f'(2)$
- (E) $f'(2) < f''(2) < f(2)$

CalcAB Sample Exam II (online version)

31. Let f be a function that is everywhere differentiable. The value of $f'(x)$ is given for several values of x in the table below.

x	-10	-5	0	5	10
$f'(x)$	-2	-1	0	1	2

If $f'(x)$ is always increasing, which statement about $f(x)$ must be true?


- (A) $f(x)$ has a relative minimum at $x = 0$.
 - (B) $f(x)$ is concave downwards for all x .
 - (C) $f(x)$ has a point of inflection at $(0, f(0))$.
 - (D) $f(x)$ passes through the origin.
 - (E) $f(x)$ is an odd function.
- Answer

32. A certain species of fish will grow from x million to $x(15 - x)$ million each year. In order to sustain a steady catch each year, a limit of $x(15 - x) - x$ million fish are to be caught, leaving x million fish to reproduce each year. What is the number of fish which should be left to reproduce each year so that the maximum catch may be sustained from year to year?

- (A) 5 million
 - (B) 7 million
 - (C) 7.5 million
 - (D) 10 million
 - (E) 15 million
- Answer

CalcAB Sample Exam II (online version)

33.




The figure above shows the graph of a function $f(x)$ which has horizontal asymptotes of $y = 3$ and $y = -3$. Which of the following statements are true?

- I. $f'(x) < 0$ for all $x \geq 0$
- II. $\lim_{x \rightarrow +\infty} f'(x) = 0$
- III. $\lim_{x \rightarrow -\infty} f'(x) = 3$

- (A) I only
- (B) II only
- (C) III only
- (D) I and II only
- (E) I, II, and III

34.




Let $g(x) = \int_a^x f(t) dt$, where $a \leq x \leq b$. The figure above shows the graph of g on $[a, b]$.

Which of the following could be the graph of $y = \frac{d}{dx} f(x)$ on $[a, b]$?

- (A)
- (B)
- (C)
- (D)
- (E)

CalcAB Sample Exam II (online version)

35.



The graph above shows the distance $s(t)$ from a reference point of a particle moving on a number line, as a function of time. Which of the points marked is closest to the point where the acceleration first becomes negative?

- (A) A
- (B) B
- (C) C
- (D) D
- (E) E

36. The derivative of f is given by $f'(x) = e^x(-x^3 + 3x) - 3$ for $0 \leq x \leq 5$.

At what value of x is $f(x)$ an absolute minimum?

- (A) For no value of x
- (B) 0
- (C) 0.618
- (D) 1.623
- (E) 5

CalcAB Sample Exam II (online version)

37.

x	$f(x)$
3.99800	1.15315
3.99900	1.15548
4.00000	1.15782
4.00100	1.16016
4.00200	1.16250

The table above gives values of a differentiable function f . What is the approximate value of $f'(4)$?

- (A) 0.00234
- (B) 0.289
- (C) 0.427
- (D) 2.340
- (E) $f'(4)$ cannot be approximated from the information given.

38. Consider the function $f(x) = \begin{cases} \frac{\sin x}{x}, & x \neq 0 \\ k, & x = 0 \end{cases}$

In order for $f(x)$ to be continuous at $x = 0$, the value of k must be

- (A) 0
- (B) 1
- (C) -1
- (D) π
- (E) a number greater than 1

CalcAB Sample Exam II (online version)

39. In the interval $0 \leq x \leq 5$ the graphs of $y = \cos 2x$ and $y = \sin 3x$ intersect four times. Let A, B, C, and D be the x -coordinates of these points so that $0 < A < B < C < D < 5$. Which of the definite integrals below represents the largest number?

(A) $\int_0^A (\cos 2x - \sin 3x) dx$

(B) $\int_A^B (\sin 3x - \cos 2x) dx$

(C) $\int_B^C (\sin 3x - \cos 2x) dx$

(D) $\int_C^D (\cos 2x - \sin 3x) dx$

(E) $\int_C^D (\sin 3x - \cos 2x) dx$

40. The function $f(x) = \tan(3^x)$ has one zero in the interval $[0, 1.4]$. The derivative at this point is

(A) 0.411

(B) 1.042

(C) 3.451

(D) 3.763

(E) undefined

CalcAB Sample Exam II (online version)

41.

x	0	1	2	3	4	5	6
$f(x)$	0	0.25	0.48	0.68	0.84	0.95	1

For the function whose values are given in the table above, $\int_0^6 f(x) dx$ is approximated by a Riemann Sum using the value at the midpoint of each of three intervals of width 2. The approximation is

- (A) 2.64
- (B) 3.64
- (C) 3.72
- (D) 3.76
- (E) 4.64

42. $\frac{d}{dx} \int_x^{x^3} \sin(t^2) dt =$

- (A) $\sin(x^6) - \sin(x^2)$
- (B) $6x^2 \sin(x^3) - 2 \sin x$
- (C) $3x^2 \sin(x^6) - \sin(x^2)$
- (D) $6x^5 \sin(x^6) - 2 \sin(x^2)$
- (E) $2x^3 \cos(x^6) - 2x \cos(x^2)$

CalcAB Sample Exam II (online version)

13. A tank is being filled with water at the rate of $300\sqrt{t}$ gallons per hour with $t > 0$ measured in hours. If the tank is originally empty, how many gallons of water are in the tank after 4 hours?

(A) 600

(B) 900

(C) 1200

(D) 1600

(E) 2400

44. The region in the first quadrant enclosed by the graphs of $y = x$ and $y = 2\sin x$ is revolved about the x -axis. The volume of the solid generated is

(A) 1.895

(B) 2.126

(C) 5.811

(D) 6.678

(E) 13.355

45. If $y = xe^x$, then $\frac{d^n y}{dx^n} =$

(A) e^x

(B) e^{nx}

(C) $(x + n)e^x$

(D) $x^n e^x$

(E) $(x + n^2)e^x$

Section II Part A: Graphing Calculator MAY BE USED.

1. Let $f(x) = a(36 - x^2)$ for $0 < a < 1$ and let R be the region in the first quadrant bounded by the y -axis, the graph of f , and the graph of $g(x) = 36 - x^2$.
 - (a) Find the area of R in terms of a .
 - (b) Find, in terms of a , the equation of the line L , tangent to the graph of f at the point $(6, 0)$.
 - (c) Determine the value of a such that the line L divides the region R into two parts with equal areas. Show the analysis that leads to your conclusion.

CalcAB Sample Exam II (online version)

Section II Part A: Graphing Calculator MAY BE USED.

2. The temperature $T(x)$, in °F, in a small office building without air conditioning is given by $T(x) = 73 - 14 \cos\left(\frac{\pi(x - 3.4)}{12}\right)$ where x is the time elapsed since midnight, $0 \leq x \leq 24$.


To cool the building, the air conditioning is turned on when the temperature first reaches the desired temperature T_0 and left on until the office closes at 6:00 p.m. (that is, when $x = 18$).

The cost per day, in dollars, of cooling is given by $C(x) = 0.16 \int_x^{18} (T(x) - T_0) dx$ for $T(x) \geq T_0$.

- Estimated to the nearest half-hour, at what time will the temperature first reach 70°F?
- Estimated to the nearest half-hour, at what time will the temperature first reach 77°F?
- What is the cost per day of cooling the office if the desired temperature is 70°F? Show your reasoning.
- How much is saved per day if the desired temperature is raised to 77°F?

Section II Part A: Graphing Calculator MAY BE USED.

3. This problem deals with functions defined by $f(x) = x^3 - 3bx$ with $b > 0$.
- In the viewing window provided below, graph the members of the family $f(x) = x^3 - 3bx$ with $b = 1$, $b = 2$, and $b = 3$. Label each graph.



Viewing Window
 $[-5, 5] \times [-15, 15]$

- Find the x - and y -coordinates of the relative maximum points of f in terms of b .
- Find the x - and y -coordinates of the relative minimum points of f in terms of b .
- Show that for all values of $b > 0$, the relative maximum and minimum points lie on a function of the form $y = -ax^3$ by finding the value of a .


CalcAB Sample Exam II (online version)

Section II Part B: Graphing Calculator MAY NOT BE USED.

5. Consider the relation defined by the equation $\tan y = x + y$ for x in the open interval $-2\pi < x < 2\pi$.
- Find $\frac{dy}{dx}$ in terms of y .
 - Find the x - and y -coordinate of each point where the tangent line to the graph is vertical.
 - Find $\frac{d^2y}{dx^2}$ in terms of y .

Section II Part B: Graphing Calculator MAY NOT BE USED.

6.



The figure above shows the graph of f' , the derivative of f . The domain of f is $-4 \leq x \leq 4$. The derivative of f is an even function and $f'(-3) = f'(3) = 0$.

- For what value of x in the interval $-4 \leq x \leq 4$ does f have a relative maximum? Justify your answer.
- For what value of x in the interval $-4 \leq x \leq 4$ does f have a relative minimum? Justify your answer.
- For what values of x is the graph of f concave downward. Use f' to justify your answer.
- If $f(0) = 0$, find the value of $\int_{-a}^a f(x) dx$. Justify your answer.