

## DiffEq - Ch 5 - Extra Practice

#1b. A mass weighing 6 pounds is attached to a spring whose spring constant is 18 lb/ft. What is the period of simple harmonic motion?

#3b. A mass weighing 10 pounds, attached to the end of a spring, stretches it 2 inches. Initially, the mass is released from rest from a point 4 inches below the equilibrium position. Find the equation of motion.

#2b. A 10-kilogram mass is attached to a spring. If the frequency of simple harmonic motion is

$$\frac{4}{\pi} \text{ cycles/s} :$$

a) What is the spring constant  $k$ ?

b) What is the frequency of simple harmonic motion if the original mass is replaced with an 80-kilogram mass?

#4b. A mass weighing 5 pounds is attached to a spring. When set in motion, the spring/mass system exhibits simple harmonic motion. Determine the equation of motion if the spring constant is  $2 \text{ lb/ft}$  and the mass is initially released from a point 8 inches below the equilibrium position with a downward velocity of  $\frac{3}{4} \text{ ft/s}$ .

#5b. A mass weighing 8 pounds is attached to a spring whose constant is  $3 \text{ lb/ft}$ . The medium offers a damping force that is numerically equal to the instantaneous velocity. The mass is initially released from a point 2 foot above the equilibrium position with a upward velocity of  $8 \text{ ft/s}$ .

- a) Determine the time at which the mass passes through the equilibrium position.
- b) Find the time at which the mass attains its extreme displacement from the equilibrium position.
- c) What is the position of the mass at this instant?

#6b. A 4-foot spring measures 12 feet long after a mass weighing 6 pounds is attached to it. The medium through which the mass moves offers a damping force numerically equal to  $\sqrt{5}$  times the instantaneous velocity.

a) Find the equation of motion if the mass is initially released from the equilibrium position

with a downward velocity of  $4 \frac{ft}{s}$ .

b) Find the time at which the mass attains its extreme displacement from the equilibrium position.

c) What is the position of the mass at this instant?

#7b. A 3-kilogram mass is attached to a spring whose constant is  $20 \frac{N}{m}$ , and the entire system is then submerged in a liquid that imparts a damping force numerically equal to 8 times the instantaneous velocity.

a) Find the equation of motion if the mass is initially released from rest from a point 2 meter above the equilibrium position.

b) Find the equation of motion if the mass is initially released from a point 2 meter below the equilibrium position with an downward velocity of  $10 \frac{m}{s}$ .

#8b. A force of 6 pounds stretches a spring 1 foot. A mass weighing 4 pounds is attached to the spring, and the system is then immersed in a medium that offer a damping force that is numerically equal to 0.5 times the instantaneous velocity.

a) Find the equation of motion if the mass is initially released from rest from a point 2 foot above the equilibrium position.

b) Use the fact that...

$$A \sin(\omega t + \phi) = C_1 \cos(\omega t) + C_2 \sin(\omega t)$$

$$\text{where } A = \sqrt{C_1^2 + C_2^2} \quad \text{and} \quad \tan \phi = \frac{C_1}{C_2}$$

...to express the equation of motion as a sum of two terms with the same frequency without phase shift.

c) Find the first time at which the mass passes through the equilibrium position heading upward.

#9b. A mass weighing 12 pounds stretches a spring  $\frac{7}{3}$  feet. The mass is initially released from rest from a point 1 foot above the equilibrium position, and the subsequent motion takes place in a medium that offers a damping force that is numerically equal to  $\frac{1}{3}$  the instantaneous velocity. Find the equation of motion if the mass is driven by an external force equal to  $f(t) = 5 \sin(5t)$ .

#10b. For the given LRC series circuit...

$$L = \frac{5}{2} H, \quad R = 8 \Omega, \quad C = \frac{1}{20} F,$$

$$E(t) = 250 V, \quad q(0) = 0 \text{ Coulombs}, \quad i(0) = 0 A$$

- a) Find the charge on the capacitor as a function of time.
- b) Find the maximum charge on the capacitor.



#11b. For the given LRC series circuit...

$$L = 2 \text{ H}, \quad R = 5 \, \Omega, \quad C = 0.5 \text{ F},$$

$$E(t) = 80 \sin(t) \text{ V}$$

- a) Find the steady-state charge on the capacitor.
- b) Find the steady-state current in the circuit.