

## 1: Definitions

### Law of Large Numbers

For a small number of trials, anything can happen. As number of trials increases, the **experimental probability** approaches the **theoretical probability**.

### Definitions

**Trial:** One complete 'occurrence' of a situation.

**Outcome:** One possible result that can occur when a trial is conducted.

**Sample space:** Set of all possible outcomes that can occur in a trial.

**Event:** Any subset of the sample space (the "desired" outcomes).

**Union:**  $A \cup B$  (A OR B)

**Intersection:**  $A \cap B$  (A AND B)

**Parameter:** Number describing a *population (or model)*, e.g.  $\mu, \sigma$

**Statistic:** Number describing a *sample*, e.g.  $\bar{x}, s$

## 2: Equally-likely outcomes

If all outcomes are equally likely:

$$P(E) = \frac{\text{number of outcomes in event } E}{\text{number of outcomes in the sample space } S}$$

$$= \frac{\text{number of 'desired' outcomes}}{\text{total number of outcomes}}$$

## 2: Counting Strategies

### Strategy

- List out all the cases and just count them up. (can also use tree diagrams, grids for 2 dice, methodical listing system to help)
- Multiplication Principle (one box per choice, fill in with number of ways to make that choice, multiply)
- Permutations (use calculator) (special case: 'choose all' =  $n!$  ways)
- Combinations (use calculator)
- Multiplication Principle w/Combinations
- Distinguishable permutations  
# distinguishable permutations =  $\frac{n!}{n_1!n_2!n_3!...}$

### When to Use


- Best strategy, but only good for small numbers.
- Multiple choice to make, every possible choice in each box can be paired with every other choice.
- A set of distinct objects (no repeats), choosing some or all, and objects are 'used up' as you choose them, and order matters.
- A set of distinct objects (no repeats), choosing some or all, and objects are 'used up' as you choose them, and order does not matter.
- Multiple choices to make, but each is a choice of a number of items out of a set.
- Number of ways to arrange all items in a set if there are repeats.

## 4: Compound Events (OR)

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

**Special Case:** If two events have no overlap, they are called 'mutually exclusive' or 'disjoint' events.

Picture this...



So the OR formula is simplified...

$$P(A \cup B) = P(A) + P(B) - \cancel{P(A \cap B)}$$



$$(P(A \cap B) = 0)$$

## 5: Compound Events (AND)

$$P(A \cap B) = P(A) \cdot P(B | A)$$

$$P(A \cap B) = P(B) \cdot P(A | B)$$

Picture a part of a tree...



**Special Case:** If the probability of an event does not change regardless of whether or not another event happens, then the events are **independent events**.

For independent events:  $P(B) = P(B | A)$

So...  $P(A \cap B) = P(A) \cdot P(B | A)$  ...simplifies to...  $P(A \cap B) = P(A) \cdot P(B)$

## 3: Conditional Probability

$$P(\text{video games} | \text{girl}) = \frac{12}{60} = .20$$

event


The event is always contained within the conditional sample space.

condition

The condition is always just a portion of the sample space (the **conditional sample space**).

The **conditional sample space** is a portion of the **sample space**.

The **event** is a portion of the **conditional sample space**.



## 3: Conditional Probability

The event goes in the numerator of the fraction.


$$P(\text{video games} | \text{girl}) = \frac{12}{60} = .20$$

The condition goes in the denominator of the fraction.

$$P(A | B) = \frac{P(A \cap B)}{P(B)}$$

$$P(\text{video games} | \text{girls}) = \frac{P(\text{video games} \cap \text{girl})}{P(\text{girl})}$$


## 4: Disjoint Events



A and B are mutually-exclusive

A and B are disjoint events

$$P(A \cap B) = 0$$



A and B are non mutually-exclusive

A and B are not disjoint events

A and B are joint events

$$P(A \cap B) \neq 0$$

## 5: Independent Events

### Test for independent events:

Two events are independent if:

$$P(B) = P(B | A) = P(B | \bar{A})$$

(check any two)

**Note:** Some books also use the simplified version of the AND formula as a 'test for independence'...


$$\text{If } P(A \cap B) = P(A) \cdot P(B)$$

then A and B are independent

...but this is more a consequence of independence, not the reason.

### 6: AND/OR together

We often need to use the AND and OR rules together:



$$B = (A \text{ and } B) \text{ or } (\bar{A} \text{ and } B)$$

$$P(B) = P(A \text{ and } B) \text{ or } P(\bar{A} \text{ and } B)$$

$$P(B) = P(A \text{ and } B) + P(\bar{A} \text{ and } B)$$

$$P(B) = (P(A) \cdot P(B|A)) + (P(\bar{A}) \cdot P(B|\bar{A}))$$


$$P(B) = (.05)(.90) + (.95)(.15)$$

$$P(B) = .1875$$

### 6: Bayes' Formula


$$P(A|E) = \frac{P(A) \cdot P(E|A)}{P(E)}$$

But use probability of paths on a tree diagram:



$$P(F|E) = \frac{(0.8)(0.4)}{(0.8)(0.4) + (0.2)(0.7)}$$

### 7: Venn Diagrams



Venn diagrams are great for word problems with lots of information.

Always start with most overlapped region, and don't forget to subtract what has already been accounted for.


You can fill with either counts or probabilities (but be consistent).

### 8: Discrete Probability Models

#### Binomial

- Only 2 outcomes
- Probabilities must be the same each trial.
- Probabilities of trials must be independent.
- Must have fixed number of trials, n

Best for: independent trials, fixed number of trials (known n), finding probability of k out of n.



$$P(\text{exactly } k \text{ successes out of } n \text{ trials}) = {}_n C_k (p)^k (q)^{n-k}$$

### 9: Expected Value vs. Average Value

$$\text{Expected Value} = \sum X \cdot P(X)$$

$$\text{Average Value} = \frac{\sum X}{n}$$


$$= \sum X \cdot \frac{1}{n}$$

(Expected Value is an average weighted by the probabilities of each outcome.)

(Average Value is an expected value for equally-likely events which each have the same probability.)


## 10: The Normal Model

Many values which have a continuous, infinite number of possible outcomes, especially quantities found in natural systems, can be modeled with a Normal distribution.



2 calculator functions for use with a Normal distribution:

**Have boundaries → Need area**




$$\text{area} = \text{normalcdf}(\text{left boundary}, \text{right boundary}, \mu, \sigma)$$

$$\text{area} = \text{normalcdf}(-999, x, \mu, \sigma)$$

Normal distributions are symmetrical, centered at a mean  $\mu$

The average distance data is from this mean (on both sides) is called the standard deviation  $\sigma$


**Have area → Need boundary**



$$\text{upper boundary} = \text{invNorm}(\text{left area}, \mu, \sigma)$$


$$x = \text{invNorm}(\text{left area}, \mu, \sigma)$$

## 10: Normal Approximation of Binomial Model




$$p = 0.2, n = 5$$

$$(np = 1)$$




$$p = 0.2, n = 10$$

$$(np = 2)$$



$$p = 0.2, n = 20$$

$$(np = 4)$$



$$p = 0.2, n = 50$$

$$(np = 10)$$

$$p = 0.2, n = 100$$

$$(np = 20)$$

Can use Normal approximation for the Binomial distribution.

If  $np \geq 10$  and  $nq \geq 10$

a Binomial distribution can be approximated with a Normal distribution with:  $\mu = np$

$$\sigma = \sqrt{npq}$$